


BMJ Open How well do mothers recall their own and their infants' perinatal events? A two-district study using cross-sectional stratified random sampling in Bihar, India

Joseph James Valadez ¹, Baburam Devkota,¹ Caroline Jeffery,¹ Wilbur C Hadden²

To cite: Valadez JJ, Devkota B, Jeffery C, *et al*. How well do mothers recall their own and their infants' perinatal events? A two-district study using cross-sectional stratified random sampling in Bihar, India. *BMJ Open* 2019;**9**:e031289. doi:10.1136/bmjopen-2019-031289

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-031289>).

Received 26 April 2019

Revised 18 November 2019

Accepted 18 November 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY. Published by BMJ.

¹Department of International Public Health, Liverpool School of Tropical Medicine, Liverpool, UK

²Department of Sociology, University of Maryland at College Park, College Park, Maryland, USA

Correspondence to

Dr Joseph James Valadez; joseph.valadez@lstmed.ac.uk

ABSTRACT

Objective Global monitoring of maternal, newborn and child health (MNCH) programmes use self-reported data subject to recall error which may lead to incorrect decisions for improving health services and wasted resources. To minimise this risk, samples of mothers of infants aged 0–2 and 3–5 months are sometimes used. We test whether a single sample of mothers of infants aged 0–5 months provides the same information.

Design An annual MNCH household survey in two districts of Bihar, India (n=6 million).

Participants Independent samples (n=475 each) of mothers of infants aged 0–5, 0–2 and 3–5 months.

Outcome measures Main analyses compare responses from the samples of infants aged 0–5 and 0–2 months with Mantel-Haenszel-Cochran statistics using 51 indicators in two districts.

Results No measurable differences are detected in 79.4% (81/102) comparisons; 20.6% (21/102) display differences for the main comparison. Subanalyses produce similar results. A difference detected for exclusive breast feeding is due to premature complementary feeding by older infants. Measurable differences are detected in 33% (8/24) of the indicators on Front Line Worker (FLW) support, 26.9% (7/26) of indicators of birth preparedness and place of birth and attendant, and 9.5% (4/42) of the indicators on neonatal and antenatal care.

Conclusions Differences in FLW visits and compliance with their advice may be due to seasonal effects: mothers of older infants aged 3–5 months were pregnant during the dry season; mothers of infants aged 0–2 months were pregnant during the monsoons, making transportation difficult. Useful coverage estimates can be obtained by sampling mothers with infants aged 0–5 months as with two samples suggesting that mothers of young infants recall their own perinatal events and those of their children. For some indicators (eg, exclusive breast feeding), it may be necessary to adjust targets. Excessive stratification wastes resources, does not improve the quality of information and increases the burden placed on data collectors and communities which can increase non-sampling error.

Strengths and limitations of this study

- The data were produced using stratified random sampling with no apparent design effect leading to an efficient use of information.
- Data were collected from female participants by female data collectors which is likely to have reduced non-sampling error.
- The large study population covers a large geographical area, reducing the likelihood that the results are pertinent only to a small group of mothers with infants and may be generalisable.
- Both weighted and unweighted results are presented giving strength to the conclusions.
- Due to insufficient overlap of variables in the 0–5 months' sample and the 3–5 months' sample, comparison between the 3–5 and the 0–5 months' sample was not possible.

INTRODUCTION

The progress towards United Nations' Sustainable Development Goal (SDG) 3 is measured with nine targets, including the maternal mortality ratio (MMR) and the under-five mortality rate (U5MR).^{1,2} In India, Bihar is one of the largest (population 110 million) and poorest (53% of households are in the lowest wealth index quintile of India³) states with high child and maternal mortality (U5MR=54, MMR=208),⁴ and is a priority for donor support for health systems strengthening (see the study by Karvande *et al*⁵ for an evaluation of the healthcare system in Bihar).

To accelerate progress towards achieving SDG 3, state governments in India pursue programmes of community-based care (see the studies by Mohan *et al* and Neogi *et al*^{6,7} for descriptions and assessments of this approach). Since 2011, the Bihar Ministry of Health has supported an Integrated Family

Health Initiative to improve the availability, quality and use of prenatal, perinatal and postnatal care for mothers and infants.⁸

The usual way to monitor progress towards achieving these goals is with household surveys. Perhaps the most commonly used surveys are cluster sample surveys such as the Demographic and Health Surveys and the Multiple Indicator Cluster Surveys.^{9 10} An alternative design is Lot Quality Assurance Sampling (LQAS) which provides comparable data but is decentralised to local health services organisations and more useful for management and programme planning.¹¹ Several states in India find it benefits their programmes.¹² Surveys rely on the reports of mothers of infants and young children, but these reports are subject to several sources of potential error and bias through interviewees not knowing, forgetting and having memory errors.^{13 14} Studies have shown both that mothers can accurately report significant facts about the birth and care of their children many years after the event,¹⁵ but also that even immediately after giving birth mothers may misreport details.^{16–18} Studies of mothers recall of their children's vaccination status concluded that due to offsetting errors of maternal reports, the resulting data accurately measured vaccination rates¹⁹; the pattern of error revealed that mothers whose children are up-to-date or nearly so tended to underestimate their child's vaccination status while mothers whose children have few vaccinations overestimate their coverage.

To improve the validity of collected data, knowledge, practice and coverage, surveys have used samples of mothers of infants aged 0–11 months or 0–5 months and children aged 6–11 months. In Bihar, local organisations departed from this convention of sampling among these three cohorts of children under 1 year of age and have been monitoring their programmes' progress by sampling five dedicated cohorts: mothers of children aged 0–2, 3–5, 6–8, 9–11 and 12–23 months with indicators focused on antenatal care, safe delivery practices, infant and young child-feeding practices, immunisation, treatment seeking and more. To avoid the possibility of maternal recall error, each of the five cohorts was asked questions particularly relevant to a child's specific age group.

In countries such as India with high maternal and child mortality rates, regular monitoring of related health service coverage is critical to reducing these rates. However, survey designs should be affordable and sustainable for local health systems; they should also produce precise, unbiased estimates.²⁰ In this study, we explore whether information is gained by sampling cohorts of children aged 0–2 and 3–5 months or whether sample sizes can be reduced by 50% by creating one sample cohort aged 0–5 months.

The research question we address is: 'Do the health service delivery coverage estimates from a sample of mothers of infants aged 0–5 months differ from those obtained from a sample of mothers of infants ages 0–2 months?' A corollary to this question is: 'Do mothers of infants 3–5 months of age display more recall error

relative to mothers of infants 0–2 months of age for antenatal, delivery or young infant health practices?' We compare district coverage estimates obtained from two independent samples of infants aged 0–2 months and 0–5 months. The implications of this study are important for health systems researchers needing results to appraise and improve their programmes.

METHODS

To answer this question, we collected information from a sample of mothers with infants aged 0–5 months and a sample of mothers with infants aged 0–2 months in two districts. This study took place within the context of a larger survey that also sampled children aged 3–5, 6–8, 9–11 and 12–23 months. These four latter samples used questionnaires with variables that either did not overlap at all or overlapped on very few indicators with the questionnaires used to interview the 0–5 and 0–2 months' samples of infants. Due to this constraint, in this study, we only use the two aforementioned groups to assess the measurement of the indicators and refer only to them for the remainder of this paper. The household sampling design we used is a stratified random sample.²¹ Within each district, the strata are administrative units of the health system which in Bihar is called a *block*. Within each block, the primary sampling unit is the Anganwadi Centre (Community Health Subcentre) Catchment Area (ACCA); 19 ACCAs are selected from each block with probability proportional to size. From each ACCA, one respondent is randomly selected from each age group under study using segmentation sampling.^{22 23} The sample of 19 mothers in each block is chosen to maximise the probability of correctly classifying a block with reference to performance targets on health-related indicators (95% reliability) while balancing the probability (10% margin of error) of incorrectly classifying a block and thereby failing to recognise either the accomplishments of local healthcare delivery systems or the local population's healthcare needs.²² For this purpose, principles of LQAS were used along with established probability tables.^{24–26}

There are 14 and 11 blocks in Gopalganj and Aurangabad ($n=6$ million), the two districts selected for this study, respectively. The total sample sizes are: (a) Gopalganj: 19×14 blocks = 266 infants aged 0–2 months and 266 infants aged 0–5 months, and (b) Aurangabad: 19×11 blocks = 209 infants aged 0–2 months and 209 infants aged 0–5 months. The 0–5 months old sample is distributed as 60% 0–2 months old and 40% 3–5 months old.

Using summary data from each of the two samples, we analyse the data with Cochran-Mantel-Haenszel (CMH)²⁷ tests for 51 dichotomous indicators (online supplementary table S1) common to the two samples. The CMH tests theoretically have a χ^2 probability distribution with 1 df. With a sufficient number of respondents or a sufficient number of blocks, the CMH test is equivalent to a conditional logistic regression (Agresti, pp114–115²⁸). In this analysis, both the number of respondents and the number

Table 1 Number of indicators by probability of a difference between the 0–2 and 0–5 months' samples for weighted and unweighted samples

Unweighted	Weighted					
	Aurangabad		Gopalganj		Total	
	≥0.05	<0.05	≥0.05	<0.05	≥0.05	<0.05
≥0.05	40	3	41	4	81	7
<0.05	0	8	1	5	1	13

of blocks only approach sufficiency. Consequently, the calculated χ^2 and probabilities must be considered as approximations of their true values.

We calculate both unweighted and weighted estimates. The unweighted estimates permit the results from smaller blocks to have equal weight vis à vis larger ones. Since the research question concerns an analysis of which age cohort is most informative, the weighted estimates may not be as useful as the unweighted ones. However, the weighted estimates provide better point estimates of the indicators at the district level. The effect of the weights on the χ^2 statistics is to increase the contribution of the larger blocks and decrease the contribution of the smaller blocks. Hence, we report both sets of results (online supplementary tables S2-S3).

The χ^2 probability distribution puts the differences between the districts on a probability scale (online supplementary table S2). To determine meaningful differences in responses between the two age cohorts, we used a probability of 0.05 as a cut-off value and considered differences with probabilities less than 0.05 to be possibly meaningful and those with larger probabilities to be likely due to sampling errors. With 102 comparisons (51 indicators weighted or unweighted), we must expect some to exceed this cut-off by chance alone. If all of the comparisons were independent, we might randomly find five differences,

but many of the indicators measure related events (eg, number of ANC visits and tetanus toxoid vaccinations) and the weighted and unweighted estimates were similar, so these indicators were not all independent, and it was not possible to calculate an expected number of differences nor was it appropriate to interpret these probabilities as measures of 'statistical significance'.

Patient and public involvement

This study does not involve patients. Also, the public was not involved in the design, conduct and reporting of the research. The public was engaged as interviewees. To ensure local engagement, we coordinated with the Bihar Ministry of Health, local implementing non-governmental organisations and our donor. We also shared the results with them and offered further dissemination of results.

RESULTS

We find a high level of agreement between the two samples (table 1). Out of 102 weighted and unweighted comparisons between the estimates from the 0–2 and 0–5 months' samples, there is no probable difference in 81 (79.4%) in both the unweighted and weighted estimates. We detect that probable differences exist for 13 comparisons (12.7%). For the remaining eight comparisons, the weighted and unweighted estimates disagree. The weighted estimates find seven differences that the unweighted estimates do not; the unweighted estimates find one difference that the weighted estimates do not find.

For different health service domains, the number of indicator comparisons varies from two (exclusive breast feeding; EBF) to 24 concerning home visits by Front Line Worker (FLW) support (table 2). The two principal FLW are Anganwadi workers and Accredited Social Health Activists (ASHA).

In the FLW support domain, 33% of comparisons have probable differences. The neonatal health domain has 20

Table 2 Number of indicator comparisons by subject domain showing a measurable difference using weighted and unweighted estimates of 0–2 and 0–5 months' samples

Health service domain	Total comparisons	No measurable difference between 0–2 and 0–5 months' results	Measurable difference between 0–2 and 0–5 months' results			Per cent indicators with different results
			Both	Unweighted only	Weighted only	
Antenatal care	22	21	0	0	1	5
Place of birth and attendant	8	6	1	0	1	25
Birth preparedness	18	13	3	0	2	28
Front Line Worker support	24	16	6	0	2	33
Maternal health	8	8	0	0	0	0
Neonatal care	20	17	1	1	1	15
Exclusive breast feeding	2	0	2	0	0	100
Totals	102	81	13	1	7	21.9



comparisons and the birth preparedness domain has 18; in these domains 15% and 28% show probable differences, respectively. The place of birth and attendant domain, and maternal health domain each have eight comparisons with 25%, or two comparisons, and 0 comparisons, respectively, showing a possible difference. The differences between the two samples cluster around home visits from FLW and behaviours associated with birth preparedness and neonatal care. Details of these differences are listed in [table 3](#).

For two indicators, both the weighted and unweighted estimates display probable differences between the 0–2 and 0–5 months samples in both districts. For indicator #52, the proportion exclusively breast feeding, the 0–5 months' cohort has the lower estimate, and indicator #24, the proportion of mothers visited by an ASHA at least once during their last pregnancy, the 0–5 months' sample gives the higher estimate, about 74%, compared with 63% in the 0–2 months' sample (online supplementary tables S2–3).

Additional analyses comparing subsamples of mothers of infants aged 0–2 months and 3–5 months from the 0–5 months' sample, the sample of mothers of infants aged 0–2 months and the subsample of infants aged 0–2 months, and the sample of mothers of infants aged 3–5 months and the subsample of infants aged 3–5 months produced similar results (online supplementary text, tables S4a–b, S5a–b and S6).

DISCUSSION

Statement of principal findings

There are no measurable differences in coverage estimates for 79.4% (81 comparisons) of the indicator comparisons between the samples of mothers with infants aged 0–2 months versus mothers of infants aged 0–5 months; 12.7% (13 comparisons) display measurable differences. The remaining 7.8% (eight comparisons) display discrepancies between the weighted and unweighted estimates.

Strengths and weaknesses of the study

The strengths of this study are that it compares estimates from two independent samples and that there are many estimates from diverse domains. The weaknesses of this study are that the data have been collected in only two districts of one state in India and in different months of a single year, and that indicators from the sample of mothers of 3–5 months old infants comparable to those of the 0–2 months old infants, using the same questionnaire, have not been collected. Supplemental analyses comparing 0–2 and 3–5 months subsamples of the 0–5 sample did not uncover evidence of bias due to the combination of these two age groups.

Strengths and weaknesses in relation to other studies

Other studies of maternal recall bias have sought a 'gold standard' to represent reality and to evaluate measures. Our study, of course, is interested in reality, but this study compares alternative measures needed to assess the Bihar health programme. It also uses a complete sample of the age grouping under study rather than just a sub-sample of

a larger age grouping. A weakness of this approach is that the analysis does not result in a formal statistical test; our conclusion is based on the weight of the evidence.

Meaning of the study

The evidence indicates that samples of the broader group yield comparable results to those of the narrower age group. It is not necessary to double the total sample by measuring independently 0–2 months' and 3–5 months' cohorts of children. These results also tend to dispel the hypothesis that maternal recall is problematic for mothers during the first 6 months following delivery. Our results are more consistent with conclusions presented in earlier research,¹⁵ and they support those organisations collecting data with 0–5 months' cohorts.

Indicator #52, EBF, displays two comparisons measuring decreases in both districts. This is not surprising as fewer infants are expected to be exclusively breast fed in a sample ranging from 0 to 5 months than a sample ranging 0–2 months; mothers introduce complementary feeding and liquids as infants age despite this being a health risk. This difference could be accommodated by adjusting expectations and targets for the indicator.

Unanswered questions and future research

Further investigation and consideration of the differences is warranted. The eight differences found in the FLW support indicators deserve more scrutiny. Seven show higher estimates for the 0–5 months' cohort, and one has a higher estimate for the 0–2 months' cohort. The former seven differences may be due to excessive rainfall during July–September (monthly 2016 average 288 mm, range: 151–35 mm) versus the lesser rainfall during October–June (monthly 2016 average 33 mm, range: 0.0–129 mm) which in the last trimester may have reduced the access of ASHA in the 0–2 months' cohort.²⁹ Indicators such as these may be particularly sensitive to rainfall and may explain why more mothers in the 0–5 months' cohort displayed higher FLW visitation estimates since FLW were not impeded by the monsoon and the resulting muddy roads.

Differences in birth preparedness and institutional birth may be a consequence of differences in rainfall or in FLW support; the results signal a need for more careful planning when transportation is difficult and decreases the effectiveness of FLW by reducing their access to women. Or, some of these differences may just be due to noise in the data.

CONCLUSIONS

Overall, the answer to the research question, 'Can one get the same district coverage estimates from a sample of mothers of infants aged 0–5 months as from a sample of mothers of infants aged 0–2 months?' is yes. This result can be paraphrased as: mothers do not display increased recall errors of their perinatal healthcare behaviour in a cohort of mothers with infants aged 0–5 months as

Table 3 Indicators by health service domain showing measurement differences between 0–2 and 0–5 months' samples

Health service domain and indicator	Indicator no	District	Weighted coverage (%)		P value		Estimate type
			0–2 months	0–5 months	Unweighted estimate	Weighted estimate	
Antenatal care							
Proportion of mothers of infants (0–2/0–5 months) registered during their last pregnancy	1	Aurangabad	85.2	77.6	0.0552	0.0365	Weighted
Place of birth and attendant							
Proportion of mothers of infants (0–2/0–5 months) whose last child was delivered at a public facility	38	Gopalganj	51.2	61.6	0.0363	0.0159	Both
Proportion of mothers of infants (0–2/0–5 months) whose last child was delivered at a health facility (private or public facility)	37	Gopalganj	78.9	85.5	0.0643	0.0459	Weighted
Birth preparedness							
Proportion of mothers of infants (0–2/0–5 months) who planned transportation to health facility in their last pregnancy (home and institutional delivery)	15	Gopalganj	45.7	56.0	0.0266	0.0158	Both
Proportion of mothers of infants (0–2/0–5 months) who identified persons to care for the baby immediately after birth (home+institutional delivery)	17	Gopalganj	51.8	62.6	0.0255	0.0103	Both
Proportion of mothers of infants (0–2/0–5 months) who planned for institutional delivery and identified person to accompany her during the delivery	23	Aurangabad	62.5	47.0	0.0039	0.0052	Both
Proportion of mothers who planned for institutional delivery of infants (0–2/0–5 months) who had a new blade and thread for their delivery	19	Aurangabad	23.5	14.5	0.062	0.0429	Weighted
Proportion of mothers who planned institutional delivery of infants (0–2/0–5 months) who arranged clean cloth for mothers and baby	21	Aurangabad	43.6	31.2	0.0546	0.0137	Weighted
FLW support							
Proportion of mothers of infants (0–2/0–5 months) who were visited by ASHA at least once during their last pregnancy	24	Aurangabad	62.2	75.2	0.0023	0.0042	Both
		Gopalganj	63.5	73.0	0.0284	0.0175	Both
Proportion of mothers of infants (0–2/0–5 months) visited at home by FLWs at least once during their last pregnancy	26	Aurangabad	63.5	76.7	0.0021	0.0032	Both
Proportion of mothers of infants (0–2/0–5 months) visited at home by ASHA within 24 hours of last delivery	31	Aurangabad	29.9	44.9	0.0009	0.0016	Both
Proportion of mothers of infants (0–2/0–5 months) visited at home by any FLW within 24 hours of last delivery	33	Aurangabad	32.2	46.7	0.0015	0.0026	Both
Proportion of mothers of infants (0–2/0–5 months) visited at home by any FLW within first week of last delivery	35	Aurangabad	44.5	59.3	0.0018	0.0026	Both
Proportion of mothers of infants (0–2/0–5 months) visited at home by any Anganwadi worker (AWW) within the first week of the last delivery	34	Gopalganj	14.4	8.9	0.0959	0.0471	Weighted
Proportion of mothers of infants (0–2/0–5 months) visited by ASHAs at least once during their last trimester of pregnancy	27	Gopalganj	52.6	61.1	0.0617	0.0449	Weighted
Infant care							
Proportion of infants aged 0–2/0–5 months who were delivered at home continued with dry cord care	51	Aurangabad	78.0	45.4	0.0001	0.0006	Both
		Gopalganj	63.7	41.1	0.0431	0.0627	Unweighted
Proportion of infants aged 0–2/0–5 months weighed after birth (public facility/private facility/home)	48	Gopalganj	70.7	78.2	0.0727	0.0464	Weighted

Continued



Table 3 Continued

Health service domain and indicator	Indicator no	District	Weighted coverage (%)		P value		Estimate type
			0–2 months	0–5 months	Unweighted estimate	Weighted estimate	
Exclusive breast feeding							
Proportion of infants (0–2/0–5 months) breast fed in the past 24 hours (exclusively breast fed)	52	Aurangabad	69.2	59.7	0.0229	0.0411	Both
		Gopalganj	82.1	68.4	0.0001	0.0003	Both

ASHA, Accredited Social Health Activist; FLW, Front Line Worker.

compared with mothers with younger infants. Substantial resources and effort can be saved using a survey design that avoids needless expenses to collect data that provides insubstantial amounts of information. It also reduces the burden on data collectors and community participants. Fatigue to both groups can result in needless non-sampling error.

Acknowledgements We gratefully acknowledge the essential roles of Hemant Das, Alok Prahdan and Sanjay Biswa for their careful field work supporting the LQAS survey and the HMIS data retrieval. We thank Professor Imelda Bates, Professor Brian Faragher and Nancy Vollmer for their valuable feedback on an earlier version of this manuscript.

Contributors JJV, BD developed the research question and survey design; WCH, CJ carried out the statistical analyses; JJV obtained the funding and donor support for the research; BD trained and managed the survey teams in Bihar; JJV, WCH, CJ interpreted the data; CJ responsible for data curation; all authors wrote and reviewed the paper.

Funding This research was funded by the Bill and Melinda Gates Foundation Investment ID OPP1142889

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The Ethical Committees of the Indian Institute of Public Health (No IIPHB-IEC-2016/010) and the Liverpool School of Tropical Medicine Research Ethics Committee approved the protocol, study instruments and consent procedures for the data collection of the household surveys (Research Protocol 16-023).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iD

Joseph James Valadez <http://orcid.org/0000-0002-6575-6592>

REFERENCES

- UNSTATS. Official list of mdg indicators: United nations statistics division, department of economic and social Affairs, 2005. Available: <https://unstats.un.org/unsd/mdg/Host.aspx?Content=Indicators/OfficialList.htm>
- UN DESA. Sustainable development goals: division for sustainable development goals, United nations department of economic and social Affairs, 2015. Available: <https://www.un.org/sustainabledevelopment/sustainable-development-goals/>
- International Institute for Population Sciences (IIPS) and ICF. *National family health survey (NFHS-4), 2015-16: India*. Mumbai: IIPS, 2017.
- MOSPI. Millennium development goals India country report 2015. New Delhi Social Statistics Division, Central Statistics Office, Ministry of Statistics and Programme Implementation; 2015.
- Karvande S, Sonawane D, Chavan S, *et al*. What does quality of care mean for maternal health providers from two vulnerable states of India? case study of Bihar and Jharkhand. *J Health Popul Nutr* 2016;35:1–10.
- Mohan P, Kishore B, Singh S, *et al*. Assessment of implementation of integrated management of neonatal and childhood illness in India. *J Health Popul Nutr* 2011;29:629–38.
- Neogi SB, Sharma J, Chauhan M, *et al*. Care of newborn in the community and at home. *J Perinatol* 2016;36:S13–17.
- CARE Family Health. Integrated family health Initiative: catalysing change for healthy communities; 2013. http://www.care.org/sites/default/files/documents/MH-2013-BIHAR_IFHI_Program%20Summary.pdf
- Blanc AK, Warren C, McCarthy KJ, *et al*. Assessing the validity of indicators of the quality of maternal and newborn health care in Kenya. *J Glob Health* 2016;6:1–11.
- Eisele TP, Rhoda DA, Cutts FT, *et al*. Measuring coverage in MNCH: total survey error and the interpretation of intervention coverage estimates from household surveys. *PLoS Med* 2013;10:e1001386.
- Anoke SC, Mwai P, Jeffery C, *et al*. Comparing two survey methods of measuring health-related indicators: lot quality assurance sampling and demographic health surveys. *Trop Med Int Health* 2015;20:1756–70.
- Valadez JJ, Devkota B, Pradhan MM, *et al*. Improving malaria treatment and prevention in India by aiding district managers to manage their programmes with local information: a trial assessing the impact of lot quality assurance sampling on programme outcomes. *Trop Med Int Health* 2014;19:1226–36.
- Tourangeau R. Remembering what happened: Memory errors and survey reports. In: Stone AA, Jobe JB, Bachrach CA, eds. *The Science of Self-report: Implications for research and practice*. Mahwah, NJ: Lawrence Erlbaum, 2000.
- Bradburn NM. Temporal representation and event dating. In: Stone AA, Jobe JB, Bachrach CA, eds. *The Science of Self-report: Implications for research and practice*. Mahwah, NJ: Lawrence Erlbaum, 2000.
- Tomeo CA, Rich-Edwards JW, Michels KB, *et al*. Reproducibility and validity of maternal recall of pregnancy-related events. *Epidemiology* 1999;10:774–6.
- Elkady E, Kenton K, White P, *et al*. Do mothers remember key events during labor? *Am J Obstet Gynecol* 2003;189:195–200.
- Li R, Scanlon KS, Serdula MK. The validity and reliability of maternal recall of breastfeeding practice. *Nutr Rev* 2005;63:103–10.
- Miles M, Ryman TK, Dietz V, *et al*. Validity of vaccination cards and parental recall to estimate vaccination coverage: a systematic review of the literature. *Vaccine* 2013;31:1560–8.
- Valadez JJ, Weld LH. Maternal recall error of child vaccination status in a developing nation. *Am J Public Health* 1992;82:120–2.
- Chan M, Kazatchkine M, Lob-Levyt J, *et al*. Meeting the demand for results and accountability: a call for action on health data from eight global health agencies. *PLoS Med* 2010;7:e1000223.
- Valadez JJ, Weiss W, Leburg C, *et al*. Assessing Community Health Programs: A Trainer's Guide. In: *Using LQAS for baseline surveys and regular monitoring*. 2nd ed. St Albans: Teaching-aids At Low Cost, 2007.
- Turner AG, Magnani RJ, Shuaib M. A not quite as quick but much cleaner alternative to the expanded programme on immunization (Epi) cluster survey design. *Int J Epidemiol* 1996;25:198–203.
- Davis RH, Valadez JJ. Improving the collection of knowledge, attitude and practice data with community surveys: a comparison of two second-stage sampling methods. *Health Policy Plan* 2014;29:1054–60.
- Dodge HF, Romig HG. *Sampling inspection tables: single and double sampling*. 2nd. New York: John Wiley & Sons, 1959.

- 25 Robertson SE, Valadez JJ. Global review of health care surveys using lot quality assurance sampling (LQAS), 1984-2004. *Soc Sci Med* 2006;63:1648-60.
- 26 Valadez JJ. *Assessing child survival programmes in developing countries: testing lot quality assurance sampling*. Boston: Harvard University Press, 1991.
- 27 Mantel N. Chi-Square tests with one degree of freedom: extensions of the Mantel-Haenszel procedure. *J Am Stat Assoc* 1963;58:690-700.
- 28 Agresti A. *An introduction to categorical data analysis*. 2nd. Hoboken, New Jersey: Wiley-Interscience; John Wiley & Sons, 2007.
- 29 Purohit, M K, and Surinder Kaur. Rainfall statistics of India -2016. New Delhi Hydromet Division, India Meteorological Department (Ministry of Earth Sciences); 2017. [hydro.imd.gov.in/hydrometweb/S\(1b2esbq4nsba2i45gfe3155\)/PRODUCTS/Publications/Rainfall%20Statistics%20of%20India%20-%202016/Rainfall%20Statistics%20of%20India%20-%202016.pdf](http://hydro.imd.gov.in/hydrometweb/S(1b2esbq4nsba2i45gfe3155)/PRODUCTS/Publications/Rainfall%20Statistics%20of%20India%20-%202016/Rainfall%20Statistics%20of%20India%20-%202016.pdf)